



## RotoSpa Granular Shock

### Features/Characteristics

- Helps prevent/remove algae
- Destroys swimmer wastes
- May be used for regular disinfection

### Application Instructions

1. Maintain pH levels within the ideal range 7.2 – 7.6. If the level is not in this range, adjust.
2. Maintain free chlorine residuals routinely in the 1 – 3mg/l (ppm) range. Shock or oxidising procedures are recommended either weekly or fortnightly with a maximum of 10mg/l (ppm) for a full shock treatment.
3. Pre-dissolve this product in a clean plastic container at a maximum rate of 100g of product of 2.2 gallon (10 litres) in water. Stir well to ensure product is fully dissolved. Add the resulting solution to the pool whilst the pump (s) are turned 'on' preferably near the water inlets to aid distribution.
4. Repeat dose/application if necessary to achieve the desired level.
5. After using this product pool water may become temporarily cloudy, so keep the filtration pump running until it clears.
6. After shock treatment, allow the chlorine to drop below 5mg/l before bathing recommences, and in any event wait for at least 30 minutes before bathing recommences.

Dosing Rates				
Water Volumes		To increase Chlorine Level		
Litres	Gallons	1 mg/l	5mg/l	10mg/l
4,545	1,000	7g	35g	70g
11,360	2,500	17g	88g	175g
22,725	5,000	35g	175g	350g
68,175	15,000	105g	525g	1050g